

中国红豆杉枝叶中的紫杉烷二萜

王福生², 彭丽艳¹, 赵 昱¹, 古 昆³, 赵勤实^{1*}, 孙汉董¹

(1 中国科学院昆明植物研究所植物化学与西部植物资源持续利用国家重点实验实, 云南 昆明 650204 ;

2 大理学院药理学系, 云南 大理 671000 ; 3 云南大学化学系, 云南 昆明 650091)

摘要: 从中国红豆杉 (*Taxus chinensis*) 枝叶的乙醇提取物中分离得到 8 个紫杉烷二萜, 通过波谱分析分别确定为: 14 β -羟基巴卡亭 VI (1), 巴卡亭 VI (2), 巴卡亭 IV (3), 1 β -去羟基巴卡亭 IV (4), 云南红豆杉酯甲 (5), 2 α -去乙酰-2 α -苯甲酰基-13 α -乙酰基云南紫杉亭 (6), 5 α -羟基-2 α , 7 β , 9 α , 10 β , 13 α -五乙酰氧基紫杉-4 (20), 11-二烯 (7) 和 Taxacin (8)。其中化合物 1 为新化合物, 并报道八个化合物在丙酮中测得的核磁共振信号, 化合物 8 的数据属首次报道。

关键词: 中国红豆杉; 红豆杉科; 紫杉烷二萜; 14 β -羟基巴卡亭 VI

中图分类号: Q 946 文献标识码: A 文章编号: 0253-2700(2003)03-0369-08

Taxoids from the Leaves and Stems of *Taxus chinensis*

WANG Fu-Sheng², PENG Li-Yan¹, ZHAO Yu¹, GU Kun³,

ZHAO Qin-Shi^{1*}, SUN Han-Dong¹

(1 State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany,

Chinese Academy of Sciences, Kunming 650204, China; 2 Pharmaceutical Department of Dali College,

Dali 671000, China; 3 Department of Chemistry, Yunnan University, Kunming 650091, China)

Abstract: A new C-14 oxygenated taxoid, 14 β -hydroxy-baccatin VI (1), together with seven known taxoids, baccatin VI (2), baccatin IV (3), 1 β -dehydroxybaccatin IV (4), taxayunnasin A (5), 2 α -deacetyl-2 α -benzoyl-13 α -acetyl taxayuntin (6), 5 α -hydroxy-2 α , 7 β , 9 α , 10 β , 13 α -pentaacetox-4 (20), 11-taxadiene (7) and taxacin (8) were isolated from the ethanolic extract of leaves and stems of *Taxus chinensis* (Pilg) Rehd. The structures of the compounds were elucidated by spectroscopic techniques. The detailed ¹³C and ¹H NMR data of the seven known compounds were measured in acetone-*d*₆, and the NMR data of taxacin (8) were reported for the first time.

Key words: *Taxus chinensis*; Taxaceae; Taxoids; 14 β -hydroxy baccatin VI

Because of the remarkable antitumor activity, report on the phytochemistry, semisynthesis, biosynthesis, and clinic use of paclitaxel and related taxoids have proliferated, and the progressing of them have been published a lot of articles (Baloglu *et al*, 1999; Kingston, 2000). Although paclita-


* To whom correspondence should be addressed. Phone (86) 8715223254. Fax: (86) 8715216343.

E-mail: qinshizhao@hotmail.com

Received date: 2003-04-14, Accepted date: 2003-04-24

作者简介: 王福生 (1965-) 男, 在职硕士研究生, 主要从事天然药物化学研究。

xel and its semisynthetic analogue have exhibited significant clinical curative effect, however, these drugs often result in a number of side effects and multidrug resistances (MDR) (Ojima *et al*, 1997). Thus, it is essential to develop the new generation of anticancer medicines, which would be possessed of superior antitumor activity and fewer side effects. In the last twenty years, many phytochemists have been devoted to isolated new taxoids and have isolated a number of new taxoids from various *Taxus species* (Li *et al*, 2002; Shinozaki *et al*, 2002; Banskota *et al*, 2002; Shen *et al*, 2002). *Taxus chinensis* (Pilg) Rehd, indigenous to China, is considered as a promising source of taxane-type diterpenoids (Zhang *et al*, 1991; Li *et al*, 1993; Tanaka *et al*, 1994). In the continuation of our research aim at new taxoids, we have further investigated on the chemical constituents of *Taxus chinensis* (Pilg) Rehd collected in Sichuan Province. As a result, a new C-14 oxygenated taxoid, 14 β -hydroxy-baccatin VI (1), together with seven known taxoids, baccatin VI (2), baccatin IV (3), 1 β -dehydroxybaccatin IV (4), taxayunansin A (5), 2 α -deacetyl-2 α -benzoyl-13 α -acetyltaxayuntin (6), 5 α -hydroxy-2 α , 7 β , 9 α , 10 β , 13 α -pentaacetox-4 (20), 11-taxadiene (7) and taxacin (8) were isolated from the ethanolic extract of the leaves and stems. The structures of the compounds were elucidated by spectroscopic techniques.




1 R₁=OH; R₂=OBz; R₃=OH

2 R₁=OH; R₂=OBz; R₃=H


3 R₁=OH; R₂=OAc; R₃=H

4 R₁=H; R₂=OAc; R₃=H




5 R₁=OAc; R₂=OH

6 R₁=OBz; R₂=OAc



7



8

Results and Discussion

Compound (1), obtained as colorless needle crystals with $[\alpha]_D^{25} + 9.42$ (c 0.57, MeOH), was determined to have formula $C_{37}H_{46}O_{15}$ by positive HRFABMS (m/z 731.2911 [$M + H$] $^+$, calcd 731.2915). The ^{13}C and DEPT NMR spectra of 1 showed 37 carbon signals which were composed of

six ester carbonyl carbons , two olefinic carbons , six aromatic ring carbons , four methyl carbons , two methylene carbons , eight methine(including seven oxygenated methine) carbons , four quaternary carbons(including two oxygenated) , and five acetyl methyl carbons , which suggested **1** had a taxoid basic skeleton , combined with the consideration of the structure of taxoids previously isolated from the genus *Taxus* plant . Furthermore , compound **1** was suggested to have a skeleton of 6/8/6 ring-system taxoid with oxetane ring deduced from the characteristic NMR signals at δ 76.8(s , C - 1) , 47.4(d , C - 3) , 46.6(s , C - 8) , 136.6(s , C - 11) , 138.7(s , C - 12) , 43.6(s , C - 15) , while the three signals at δ 82.0(C - 4) , 84.2(C - 5) , and 76.4(C - 20) corresponding to the three carbons signals of oxetane ring .

Table 1 NMR data of the compounds **1** and **2**^a

Position	1		HMBC	2	
	δ_{C}	δ_{H}		δ_{C}	δ_{H}
1	76.8 s		H-2 , 3 , 14 , 16 , 17 , OH	78.2 s	
2	73.3 d	6.03(1H , d , 6.1)	H-3 , 14	73.9 d	5.92(1H , d , 6.0)
3	47.4 d	3.18(1H , d , 6.1)	H-2 , 5 , 20 α , 20 β	48.0 d	3.24(1H , d , 6.0)
4	82.0 s		H-3 , 5 , 6 α , 6 β , 20 α	82.0 s	
5	84.2 d	4.92(1H , d , 8.9)	H-3 , 6 α , 6 β , 20 β	84.2 d	4.94(1H , d , 7.9)
6	35.4 t	2.41(1H , m)	H-7	35.3 t	2.46(1H , m)
		1.79(1H , m)			1.84(1H , m)
7	72.5 d	5.56(1H , dd , 9.8 , 8.0)	H-3 , 5 , 6 α , 6 β , 9 , 19	72.5 d	5.53(1H , dd , 9.7 , 7.8)
8	46.6 s		H-2 , 3 , 7 , 9 , 19	46.5 s	
9	75.4 d	6.14(1H , brs) ^b	H-3 , 7 , 10 , 19	75.4 d	5.97(1H , d , 11.3)
10	71.3 d	6.14(1H , brs) ^b	H-9	71.2 d	6.17(1H , d , 11.3)
11	136.6 s		H-9 , 10 , 13 , 16 , 17 , 18	139.6 s	
12	138.7 s		H-10 , 13 , 18	141.3 s	
13	79.2 d	6.07(1H , brd , 6.7)	H-14	70.2 d	6.14(1H , t , 8.7)
14	70.7 d	4.28(1H , d , 6.4)	H-2 , 13 , OH-14	36.8 t	2.37(2H , m)
		4.41(1H , s , OH-14)			
15	43.6 s		H-10 , 14 , 2 , 16 , 17	43.8 s	
16	28.7 q	1.17(3H , s)	H-17	28.4 q	1.20(3H , s)
17	24.5 q	1.74(3H , s)	H-16	23.4 q	1.75(3H , s)
18	14.5 q	1.98(3H , s)		15.2 q	1.98(3H , d , 1.0)
19	13.0 q	1.62(3H , s)	H-3 , 7 , 9	13.1 q	1.60(3H , s)
20	76.4 t	4.17(1H , d , 8.0)	H-3	76.6 t	4.18(1H , d , 7.9)
		4.12(1H , d , 8.0)			4.07(1H , d , 7.9)
OCOPh	166.1 s		H-2 , 2' , 6'	166.3 s	
i	131.0 s		H-2' , 3' , 6'	130.7 s	
o	130.7 d	8.11(2H , d , 8.3)	H-3' , 4' , 5'	131.0 d	8.09(2H , dd , 8.5 , 1.4)
m	129.4 d	7.51(2H , t , 7.8)	H-2' , 4' , 5'	129.3 d	7.52(2H , dd , 8.5 , 7.4)
p	134.1 d	7.63(1H , t , 7.4)	H-2' , 3' , 5' , 6'	134.1 d	7.62(1H , t , 7.4)
OAc	171.3 s			171.1 s	
OAc	170.9 s		H-13 , -COCH ₃	170.8 s	
OAc	170.8 s		H-9 , -COCH ₃	170.3 s	
OAc	170.3 s		H-7 , -COCH ₃	170.2 s	
OAc	169.4 s		H-10 , -COCH ₃	169.4 s	
OAc	22.8 q	2.37(3H , s)		22.9 q	2.31(3H , s)
OAc	21.4 q	2.21(3H , s)		21.4 q	2.18(3H , s)
OAc	21.1 q	2.13(3H , s)		21.2 q	2.11(3H , s)
OAc	20.8 q	2.08(3H , s)		20.9 q	2.06(3H , s)
OAc	20.8 q	1.98(3H , s)		20.8 q	1.96(3H , s)

^a ¹³C , ¹H NMR and HMBC spectra were measured at 100 , 400 and 500 MHz , respectively , in acetone-*d*₆ (δ in ppm , *J* in Hz).

^b H-9 and H-10 appeared as broad singlet in acetone-*d*₆ , while they appeared as 6.60(1H , d , 11.2 , H-9) ; 6.67(1H , d , 11.2 , H-10) in C₅D₅N .




Fig. 1 Key NOESY correlations for **1**

Comparison of the ^{13}C and ^1H NMR spectra (Table 1) of **1** with that of baccatin VI (**2**), a known taxoid isolated from the same plant this time, revealed that the methylene signals [δ_{C} 36.8 (t, C-14) and δ_{H} 2.37 (2H, m, H_2 -14)] in **2** were replaced by the methine signals [δ_{C} 70.7 (t, C-14) and δ_{H} 4.28 (1H, d, 6.4, H-14)] in **1**, which was approved by the ^1H - ^1H COSY correlations of a methine proton (δ_{H} 4.28, H-14) with another methine proton (δ_{H} 6.07, H-13), and by the HMBC correlations of the proton signal (H-14) with carbon signals [δ_{C} 76.8 (C-1), 73.3 (C-2), 79.2 (C-13), 43.6 (C-15)]. Therefore, compound **1** differed from baccatin VI (**2**) by the presence of a hydroxyl group at C-14 in **1**, which was further confirmed by the HMBC correlation of a hydroxyl proton signal (δ_{H} 4.41, s, OH-14) with the signal of C-14. Finally, the relative stereochemistry of **1** was established by its NOESY spectrum (Figure 1). The protons H-14/H-3 and H-14/H-20 α showed correlations with each other, which indicated H-14 has α orientation. On the basis of the spectral evidence described above, the structure of **1** was established as 14 β -hydroxy baccatin VI.

The seven known compounds were identified as baccatin VI (**2**) (Senilh *et al*, 1984; Della Casa de Marcano *et al*, 1975), baccatin IV (**3**) (Della Casa de Marcano *et al*, 1975), 1 β -dehydroxybaccatin IV (**4**) (Della Casa de Marcano *et al*, 1975), taxayunnasin A (**5**) (Liu *et al*, 1992), 2 α -deacetyl-2 α -benzoyl-13 α -acetyl taxayuntin (**6**) (Chen *et al*, 1993), 5 α -hydroxy-2 α , 7 β , 9 α , 10 β , 13 α -pentaacetoxy-4 (20), 11-taxadiene (**7**) (Kingston *et al*, 1982) and taxacin (**8**) (Yoshizaki *et al*, 1988). The structures of the known compounds were elucidated by spectroscopic techniques and confirmed by comparison of spectral data with literature report.

Experimental

General The Melting point was determined on an XRC-1 micro melting point apparatus and uncorrected. NMR spectra were performed on a Bruker AM-400 MHz and DRX-500 MHz spectrometer. FABMS and HRFABMS were taken on a VG Auto Spec-3000 or on a Finnigan MAT 90 instrument. IR spectra were recorded on a Bio-Rad FTS-135 spectrometer with KBr pellets. UV spectrum was obtained on a UV 2401 PC spectrometer. Optical rotations were measured with a HORIBA SEPA-300 High Sensitive Polarimeter. Column chromatography was performed either on 200-300 mesh silica gel and 10-40 μm silica gel H; 43-63 μm Lichroprep RP-18 and Sephadex LH-20 were used for column chromatography. Fractions were monitored by TLC and spots were visualized by heating silica gel plates sprayed with 10% H_2SO_4 in EtOH.

Plant material The leaves and stems of *Taxus chinensis* (Pilg) Rehd were collected in Sichuan Province. A voucher specimen has been deposited at the Yunnan Academy of Forestry, Kunming, Yunnan, People's Republic of China.

Extraction and isolation The dried plant material (15 kg) was extracted three times with 95% ethanol to yield a

crude extract. After evaporation of the solvent, the residue was dissolved with MeOH/H₂O (9:1), and the MeOH-soluble part was further reextracted with chloroform to give the extract. The chloroform extract was chromatographed on Silica gel columns using solvents of increasing polarity (petroleum-EtOAc, 9:1 – 2:8, acetone, v/v) to give ten fractions, and three of these fractions (4 – 6, 16.65 g) was further isolated by repeated column chromatography on silica gel to give baccatin VI (0.2 g), baccatin IV (0.4 g), 1 β -dehydroxybaccatin IV (20 mg), and taxayunnansin A (1.0 g), 1 β -dehydroxy baccatin IV (15 mg), 14 β -hydroxy baccatin VI (16 mg), taxacin (12 mg) and 5 α -hydroxy-2 α , 7 β , 9 α , 10 β , 13 α -pentaacetoxo-4 (20), 11-taxadiene (95 mg).

14 β -hydroxy baccatin VI (1), C₃₇H₄₆O₁₅, colorless needle crystals (acetone), mp 241 – 243°C, [α]_D²⁵ + 9.42 (c 0.57, MeOH). UV $\lambda_{\max}^{\text{MeOH}}$ (log ϵ) 228.4 (4.22), 200.2 (3.84), 274.4 (2.94) nm. IR ν_{\max}^{KBr} 3443 (OH), 1740 (ester C=O), 1636, 1437, 1374, 1250, 1106, 713 cm⁻¹. ¹H and ¹³C NMR see table 1. Positive FABMS m/z (%): 731 ([M + H]⁺, 32), 713 (53), 671 (100), 447 (13), 105 (49), 83 (37); positive HRFABMS m/z 731.2911 [M + H]⁺ (calcd for C₃₇H₄₇O₁₅, 731.2915).

Baccatin VI (2), C₃₇H₄₆O₁₄, white powder, positive FABMS m/z (%): 715 ([M + H]⁺, 25), 655 (100), 553 (10), 371 (6), 311 (3). [α]_D²⁷ - 10.93 (c 1.04, acetone). ¹H and ¹³C NMR see table 1.

Baccatin IV (3), C₃₂H₄₄O₁₄, white powder, positive FABMS m/z (%): 653 ([M + H]⁺, 18), 593 (100), 533 (13), 491 (11), 371 (6). ¹³C NMR (100 MHz, acetone-*d*₆): δ 77.7 (s, C-1), 70.2 (d, C-2), 47.9 (d, C-3), 81.8 (s, C-4), 84.3 (d, C-5), 35.3 (t, C-6), 72.5 (d, C-7), 46.4 (s, C-8), 73.3 (d, C-9), 71.3 (d, C-10), 135.0 (s, C-11), 141.3 (s, C-12), 75.6 (d, C-13), 36.9 (t, C-14), 43.7 (s, C-15), 28.4 (q, C-16), 23.2 (q, C-17), 15.1 (q, C-18), 13.1 (q, C-19), 76.6 (t, C-20); OAc: 171.1 (2C, s), 170.8 (s), 170.4 (s), 170.2 (s), 169.4 (s), 23.0 (q), 21.4 (q), 21.2 (q), 20.8 (3C, q). ¹H NMR (400 MHz acetone-*d*₆): δ 5.65 (1H, d, 5.7, H-2), 3.01 (1H, d, 5.7, H-3), 4.92 (1H, d, 8.1, H-5), 1.73 (1H, m, H-6 α), 2.40 (1H, m, H-6 β), 5.48 (1H, dd, 9.7, 7.8, H-7), 6.00 (1H, d, 11.3, H-9), 6.08 (1H, d, 11.3, H-10), 6.09 (1H, m, H-13), 2.18 (2H, overlap, H₂-14), 1.17 (3H, s, Me-16), 1.66 (3H, s, Me-17), 1.94 (3H, s, Me-18), 1.53 (3H, s, Me-19), 4.47 (1H, d, 7.8, H-20 α), 4.14 (1H, d, 7.8, H-20 β), OAc: 2.19 (3H, s), 2.14 (3H, s), 2.09 (3H, s), 2.08 (3H, s), 2.01 (3H, s), 1.98 (3H, s).

1 β -dehydroxybaccatin IV (4), C₃₂H₄₄O₁₃, colorless prisms, positive FABMS m/z (%): 637 ([M + H]⁺, 19), 577 (100), 517 (2), 475 (9). ¹³C NMR (100 MHz, acetone-*d*₆): δ 45.2 (d, C-1), 69.4 (d, C-2), 47.7 (d, C-3), 81.5 (s, C-4), 84.0 (d, C-5), 35.5 (t, C-6), 71.6 (d, C-7), 46.3 (s, C-8), 72.7 (d, C-9), 71.2 (d, C-10), 134.2 (s, C-11), 139.4 (s, C-12), 75.7 (d, C-13), 26.9 (t, C-14), 38.7 (s, C-15), 31.4 (q, C-16), 27.4 (q, C-17), 15.1 (q, C-18), 13.1 (q, C-19), 77.0 (t, C-20); OAc: 171.1 (s), 170.9 (s), 170.3 (s), 170.2 (s), 170.0 (s), 169.3 (s), 22.9 (q), 21.4 (2C, q), 21.2 (q), 20.8 (2C, q). ¹H NMR (400 MHz acetone-*d*₆): δ 2.89 (1H, d, 5.8, H-1), 5.60 (1H, dd, 5.8, 2.2, H-2), 3.2 (1H, d, 5.4, H-3), 4.98 (1H, d, 9.0, H-5), 2.43 (1H, m, H-6 α), 1.80 (1H, m, H-6 β), 5.55 (1H, dd, 9.8, 7.8, H-7), 5.95 (1H, d, 11.3, H-9), 6.10 (1H, d, 11.3, H-10), 5.92 (1H, dd, 2.9, 1.5, H-13), 1.67 (1H, m, H-14a), 2.48 (1H, m, H-14b), 1.78 (3H, s, Me-16), 1.13 (3H, s, Me-17), 1.99 (3H, s, Me-18), 1.55 (3H, s, Me-19), 4.52 (1H, d, 8.1, H-20 α), 4.20 (1H, d, 8.1, H-20 β); OAc: 2.23 (3H, s), 2.18 (3H, s), 2.12 (3H, s), 2.07 (3H, s), 2.03 (3H, s), 1.97 (3H, s).

Taxayunnansin A (5), C₃₅H₄₄O₁₃, colorless prism. ¹³C NMR (100 MHz, acetone-*d*₆): δ 68.1 (s, C-1), 71.4 (d, C-2), 45.4 (d, C-3), 79.4 (s, C-4), 82.2 (d, C-5), 35.7 (t, C-6), 69.9 (d, C-7),

44.3(s, C-8), 76.7(d, C-9), 77.6(d, C-10), 134.5(s, C-11), 152.5(s, C-12), 68.8(d, C-13), 39.9(t, C-14), 76.0(s, C-15), 26.6(q, C-16), 28.3(q, C-17), 12.0(q, C-18), 12.9(q, C-19), 74.8(t, C-20); 10-OBz: 165.0(s), 130.8(s), 130.2(2C, d), 129.5(2C, d), 134.1(d); OAc: 171.5(s), 170.5(s), 170.3(s), 169.9(s), 22.1(q), 21.8(q), 21.4(q), 20.8(q). ^1H NMR(400 MHz acetone- d_6): δ 6.18(1H, d, 7.8, H-2), 3.08(1H, d, 7.8, H-3), 4.95(1H, d, 8.5, H-5), 1.78(1H, dd, 14.9, 7.8, H-6 α), 1.73(1H, d, 6.6, H-6 β), 5.55(1H, t, 8.5, H-7), 6.24(1H, d, 10.6, H-9), 6.41(1H, d, 10.6, H-10), 4.56(1H, t, 6.8, H-13), 2.50(1H, dd, 15.1, 8.6, H-14), 2.26(1H, dd, 14.6, 7.3, H-14), 1.18(3H, s, Me-16), 1.65(3H, s, Me-17), 1.71(3H, s, Me-18), 1.14(3H, s, Me-19), 4.53(1H, d, 7.3, H-20 α), 4.18(1H, d, 7.3, H-20 β); 10-OBz: 7.89(2H, d, 8.6, H-2', 6'), 7.47(2H, t, 7.6, H-3', 5'), 7.58(1H, t, 7.6, H-5'); OAc: 2.15(3H, s), 2.12(3H, s), 2.11(3H, s), 2.02(3H, s).

2 α -deacetyl-2 α -benzoyl-13 α -acetyl-taxayuntin (6), $\text{C}_{42}\text{H}_{48}\text{O}_{14}$, amorphous powder, positive FABMS m/z (%): 777([M+H] $^+$, 15), 655(100), 537(8). ^{13}C NMR(100 MHz, acetone- d_6): δ 68.8(s, C-1), 69.1(d, C-2), 45.7(d, C-3), 79.6(s, C-4), 85.0(d, C-5), 35.7(t, C-6), 71.3(d, C-7), 44.3(s, C-8), 77.3(d, C-9), 68.9(d, C-10), 137.6(s, C-11), 148.1(s, C-12), 79.1(d, C-13), 37.2(t, C-14), 75.8(s, C-15), 27.0(q, C-16), 28.2(q, C-17), 12.0(q, C-18), 13.0(q, C-19), 74.8(t, C-20); 10-OBz: 165.1(s), 130.5(s), 130.3(2C, d), 129.5(2C, d), 134.1(d); 2-OBz: 166.8(s), 130.7(s), 130.5(2C, d), 129.6(2C, d), 134.4(d); OAc: 170.9(s), 170.3(s), 170.2(s), 170.0(s), 22.2(q), 21.4(q), 21.1(q), 20.8(q). ^1H NMR(400 MHz, acetone- d_6): δ 6.52(1H, d, 7.8, H-2), 3.17(1H, d, 7.8, H-3), 4.97(1H, d, 8.1, H-5), 2.51(1H, t, 8.6, H-6 α), 1.78(1H, t, 8.8, H-6 β), 5.60(1H, t, 8.5, H-7), 6.24(1H, d, 10.6, H-9), 6.41(1H, d, 10.6, H-10), 5.70(1H, t, 7.5, H-13), 2.01(1H, overlap, H-14 α), 2.57(1H, m, H-14 β), 1.12(3H, s, Me-16), 1.23(3H, s, Me-17), 1.87(3H, s, Me-18), 1.72(3H, s, Me-19), 4.37(1H, d, 7.7, H-20 α), 4.01(1H, d, 7.5, H-20 β), 10-OBz: 8.09(2H, d, 8.4), 7.49(2H, t, 7.9), 7.57(1H, t, 7.9); 2-OBz: 7.93(2H, d, 8.4), 7.47(2H, t, 7.9), 7.55(1H, t, 7.9), OAc: 2.27(3H, s), 2.12(3H, s), 2.06(3H, s), 1.73(3H, s).

5 α -hydroxy-2 α , 7 β , 9 α , 10 β , 13 α -pentaacetoxy-4(20), 11-taxadiene(7), $\text{C}_{30}\text{H}_{42}\text{O}_{11}$, colourless prism, EIMS(70 eV) m/z (%): 578[M] $^+$ (51), 560(40), 518(27), 458(47), 416(53), 398(62), 374(29), 356(72), 338(74), 314(34), 297(48), 278(100), 263(69), 235(43), 209(29), 163(40), 145(52), 133(70), 121(48), 105(42), 91(32), 60(30). ^{13}C NMR(100 MHz, CDCl_3): δ 48.1(d, C-1), 69.5(d, C-2), 40.6(d, C-3), 144.7(s, C-4), 75.3(d, C-5), 37.1(t, C-6), 69.5(d, C-7), 47.8(s, C-8), 75.8(d, C-9), 71.9(d, C-10), 133.9(s, C-11), 138.3(s, C-12), 70.3(d, C-13), 28.6(t, C-14), 37.2(s, C-15), 26.0(q, C-16), 32.0(q, C-17), 15.9(q, C-18), 12.9(q, C-19), 116.6(t, C-20); OAc: 170.1(s), 169.8(s), 169.6(s), 169.3(s), 169.2(s), 21.5(q), 21.4(q), 21.4(q), 21.0(q), 20.8(q). ^1H NMR(400 MHz, CDCl_3): δ 1.85(1H, d, 9.2, H-1), 5.49(1H, dd, 6.2, 1.8, H-2), 3.47(1H, d, 6.0, H-3), 4.23(1H, t, 2.8, H-5), 2.18(1H, m, H-6 α), 1.45(1H, dd, 15.8, 5.4, H-6 β), 5.60(1H, dd, 10.7, 5.2, H-7), 5.84(1H, d, 10.9, H-9), 6.20(1H, d, 10.9, H-10), 5.75(1H, m, H-13), 2.62(1H, m, H-14 α), 1.57(1H, m, H-14 β), 1.69(3H, s, Me-16), 1.00(3H, s, Me-17), 2.21(3H, s, Me-18), 0.93(3H, s, Me-19), 4.83(1H, brs, H-20a), 5.27(1H, brs, H-20b), OAc: 2.09(3H, s); 2.02(6H, s), 1.99(3H, s), 1.94(3H, s).

Taxacin (8), $C_{44}H_{48}O_{15}$, colourless lameller, positive FABMS m/z (%): 817([M + H]⁺, 9), 669(100), 609(21), 549(4), 105(100). ¹³C NMR (100 MHz, acetone- d_6): δ 48.7(s, C-1), 70.8(d, C-2), 41.4(d, C-3), 141.8(s, C-4), 75.2(d, C-5), 37.0(t, C-6), 69.5(d, C-7), 50.2(s, C-8), 69.8(d, C-9), 64.9(d, C-10), 81.4(s, C-11), 92.1(s, C-12), 205.1(s, C-13), 34.7(t, C-14), 50.0(s, C-15), 16.1(q, C-16), 82.3(t, C-17), 12.9(q, C-18), 62.2(t, C-19), 116.1(t, C-20); 19-OBz: 167.1(s), 130.6(s), 130.7(2C, d), 129.3(2C, d), 134.1(d); 5-OCinnamoyl: 165.1(s), 131.1(d), 146.1(d), 135.8(s), 130.9(2C, d), 129.6(2C, d), 131.1(d); OAc: 171.6(s), 170.1(s), 169.3(s), 168.9(s), 21.3(q), 21.2(q), 20.9(q), 20.8(q). ¹H NMR (400 MHz acetone- d_6): δ 2.68(1H, m, H-1), 6.24(1H, dd, 10.4, 2.8, H-2), 3.18(1H, d, 10.2, H-3), 5.53(1H, dd, 6.2, 2.7, H-5), 2.26(1H, ddd, 2.3, 6.4, 14.7, H-6 α), 1.84(1H, m, H-6 β), 5.56(1H, t, 6.5, H-7), 5.67(1H, d, 2.8, H-9), 5.39(1H, d, 2.8, H-10), 2.56(1H, d, 19.9, H-14 α), 3.11(1H, dd, 11.9, 19.2, H-14 β), 1.35(3H, s, Me-16), 3.96(1H, d, 7.9, H-17a), 3.66(1H, d, 7.9, H-17b), 1.12(3H, s, Me-18), 5.00(1H, d, 12.3, H-19a), 4.49(1H, d, 12.3, H-19b), 5.60(1H, s, H-20a), 4.82(1H, s, H-20b); 19-OBz: 8.21(2H, dd, 6.3, 8.2, H-3''), 7.42(2H, t, 7.4, H-3', 5'), 7.66(1H, t, 8.6, H-5''); 5-OCinnamoyl: 7.92(2H, t, 8.9, H-2', 6'), 7.42(2H, t, 7.4, H-3', 5'), 7.47(1H, m, H-4'), 7.96(1H, d, 16.2, H-7'), 6.77(1H, d, 16.2, H-8''); OAc: 2.10(3H, s); 2.07(3H, s); 2.03(3H, s); 1.92(3H, s).

Acknowledgements: The authors are grateful to all members of the Analytical Group in State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, for measurements of all spectra. This project was supported by the Special Supported Bioscience and Biotechnology Foundation of Academic Sinica (STZ-01-15).

References

- Baloglu E, Kingston DGI, 1999. The taxane diterpenoids [J]. *J Nat Prod*, **62**: 1448—1472
- Banskota AH, Usia T, Tezuka Y, *et al*, 2002. Three new C-14 oxygenated taxanes from the wood of *Taxus yunnanensis* [J]. *J Nat Prod*, **65**: 1700—1702
- Chen WM, Zhang PL, Zhao JY, *et al*, 1993. Taxayuntin A, B, C, and D—from new tetraacyclic taxanes from *Taxus yunnanensis* [J]. *Chin Chem Lett*, **4**(8): 695—698
- Della Casa de Marcano DP, Halsall TG, 1975. Structures of some taxane diterpenoids, baccatins-III, -IV, -VI, and -VII and 1-dehydroxybaccatin-IV, possessing an oxetane ring [J]. *J Chem Soc Chem Commun*, 365—366
- Kingston DGI, 2000. Recent advances in the chemistry of taxol [J]. *J Nat Prod*, **63**: 726—734
- Kingston DGI, Hawkins DR, Ovington L, 1982. New taxanes from *Taxus brevifolia* [J]. *J Nat Prod*, **45**: 466—470
- Li B, Tanaka K, Fuji K, *et al*, 1993. Three new diterpenoids from *Taxus chinensis* [J]. *Chem Pharm Bull*, **41**(9): 1672—1673
- Li SH, Zhang HJ, Yao P, *et al*, 2001. Taxane diterpenoids from the bark of *Taxus yunnanensis* [J]. *Phytochemistry*, **58**: 369—374
- Liu XK (刘锡葵), Wu DG (吴大刚), Wang ZY (王宗玉), 1992. New diterpenoids from *Taxus yunnanensis* [J]. *Chinese Science Bulletin (科学通报)*, **23**: 2186—2189
- Ojima I, Slater JC, Michaud E, *et al*, 1997. Synthesis and structure-activity relationships of nonaromatic taxoids: effects of alkyl and alkenyl ester groups on cytotoxicity [J]. *J Med Chem*, **39**: 279
- Shen YC, Wang SS, Pan YL, *et al*, 2002. New taxane diterpenoids from the leaves and twigs of *Taxus sumatrana* [J]. *J Nat Prod*, **65**: 1848—1852
- Senilh V, Bleichert S, Colin M, *et al*, 1984. Mise en Evidence de Nouveaux Analogues du Taxol Extraits de *Taxus baccata* [J]. *J*

Nat Prod , **47** : 131—137

Shinozaki Y , Fukamiya N , Fukushima M , *et al* , 2002 . Dantaxusins C and D , two novel taxoids from *Taxus yunnanensis* [J] . *J Nat Prod* , **65** : 371—374

Tanaka K , Fuji K , Yokoi T , *et al* , 1994 . On the structures of six new diterpenoids , taxchinins E , H , I , J , K and taxchin B [J] . *Chem Pharm Bull* , **42** (7) : 1539—1541

Yoshizaki F , Fuduka M , Hisamichi S , *et al* , 1988 . Structures of taxane diterpenoids from the seeds of Japanese Yew , *Taxus cuspidata* [J] . *Chem Pharm Bull* , **36** : 2098—2102

Zhang ZP , Jia ZJ , 1991 . Taxanes from *Taxus chinensis* [J] . *Phytochemistry* , **30** (7) : 2345—2348

* * * * *

《云南植物研究》启事

为了提高我们的服务质量，更好的为广大作者、读者服务。自 2003 年起，在本刊编辑部的网页上，均可查询《云南植物研究》的征稿简则、编辑部的有关通知、以及来稿的流通情况，方便作者及时查阅稿件的处理情况，缩短我们与作者交流的距离。打开本刊网页后，点击“本刊动态”或“稿约讯息”，即可进行“在审稿件题目”，“作者修改稿件题目”，“已待编稿件题目”的查询。读者可以浏览本刊最新一期的目录，并可及时征订。我们衷心希望广大作者、读者能对我们的工作提出更多的宝贵意见，以便我们及时的改进工作，更好的为大家服务。刊物的良好发展，离不开广大读者、作者的大力支持，出版一份优秀的期刊，是我们共同的愿望。对我们刊物若有好的建议和要求，请及时通知我们。我们的网址为：<http://yoke.chinajournal.net.cn>

第 25 卷 第 3 期

Vol. 25 No. 3